KETOMED™ (ketoprofen)





KetoMed is recommended for the alleviation of inflammation and pain associated with musculoskeletal disorders in horses.

The product is an FDA-approved non-narcotic, non-steroidal anti-inflammatory agent with analgesic and antipyretic properties.

As part of the trusted equine pain and sedation portfolio from Bimeda, KetoMed is safe and effective against equine pain and inflammation.

LIST No UNIT PACKAGE CASE SIZE 1KET007 50 ml 12 1KET008 100 ml 12

- ¹ Owens JG, Kamerling SG, Stanton SR, Keowen ML. Effects of ketoprofen and phenylbutazone on chronic hoof pain and lameness in the horse. *Equine Vet J.* 1995;27(4):296-300.
- 2 Owens JG, Kamerling SG, Barker SA. Pharmacokinetics of ketoprofen in healthy horses and those with acute synovitis. \textit{J Vet Pharmacol Ther.} 1995;18(3):187-195.
- ³ Longo E, Autefage A, Bayle R, Keister M, Van Gool E. Efficacy of a nonsteroidal anti-inflammatory, ketofen 10% (ketoprofen) in the treatment of colic in horses. *J Equine Vet Sci.* 1992;12(5):311-315.

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

See reverse for Administration & Dosage

BENEFITS

- ✓ Bioequivalent to Ketofen® Sterile Solution
- ✓ Bottle is shatterproof plastic
- Significantly reduces hoof pain associated with chronic laminitis for up to six hours¹
- ✓ Showed significant intra-articular anti-inflammatory activity in an experimentally induced acute carpal synovitis model²
- ✓ Therapeutically equivalent to flunixin meglumine in the alleviation of pain in carpal model of arthritis³
- ✓ NSAID of the propionic acid class that includes ibuprofen, naproxen and fenoprofen





KetoNedTM (ketoprofen)



Sterile Solution, 100 mg/mL

For intravenous use in horses only.

CAUTION

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION


Ketoprofen is a non-steroidal anti-inflammatory agent of the propionic acid class that includes ibuprofen, naproxen and fenoprofen. Each mL of KetoMed (ketoprofen) contains 100 mg of ketoprofen in an aqueous formulation containing: L-Arginine, 70 mg; citric acid (to adjust pH); benzyl alcohol, 0.025 g (as preservative).

It is packaged in a multiple dose bottle.


PHARMACOLOGY

KetoMed is a non-narcotic, non-steroidal anti-inflammatory agent with analgesic and antipyretic properties.

In horses, intravenous dosages of ketoprofen ranging from 0.5 to 1.5 mg/lb resulted in dosage dependent anti-inflammatory effects in the chronic adjuvant carpitis model as depicted in the following graph.



Additional studies using the same model in horses have shown that the effects of ketoprofen are maximal by 12 hours and still measurable at 24 hours after each dosage as depicted in the following graph.



n = number of animals

TAKE TIME OBSERVE LABEL DIRECTIONS

TOXICITY

Horses were found to tolerate ketoprofen given intravenously at dosages of 0,1,3 and 5 mg/lb once daily for 15 consecutive days (up to five times the recommended dosage for three times the usual duration) with no evidence of toxic effects. In clinical studies, intravenous injection of 1 mg/lb/day for five days resulted in no injection site irritation or other side effects.

At 15-fold overdose (15 mg/lb/day) for five days one of two horses developed severe laminitis, but no gross lesions or histologic changes were observed. The toxic effects observed in the horses given a 25-fold overdose (25 mg/lb/day) for five days included inappetence, depression, icterus, abdominal swelling and postmortem findings of gastritis, nephritis and hepatitis.

INDICATION

KetoMed (ketoprofen) is recommended for the alleviation of inflammation and pain associated with musculoskeletal disorders in the horse.

ADMINISTRATION AND DOSAGE

The recommended dosage is 1 mg/lb (1 mL/100 lbs) of body weight once daily. Treatment is administered by intravenous injection and may be repeated for up to five days. Onset of activity is within two hours with peak response by 12 hours.

CONTRAINDICATIONS

There are no known contraindications to this drug when used as directed. Intra-arterial injection should be avoided. Do not use in a horse if it has previously shown hypersensitivity to ketoprofen.

CAUTIONS

This product should not be used in breeding animals since the effects of KetoMed on fertility, pregnancy or fetal health in horses have not been determined.

PRECAUTIONS

Studies to determine activity of KetoMed when administered concomitantly with other drugs have not been conducted. Drug compatibility should be monitored closely in patients requiring adjunctive therapy.

WARNING

Do not use in horses intended for human consumption.

SIDE EFFECTS

During investigational studies, no significant side effects were reported.

To report suspected adverse drug events, for technical assistance, or to obtain a copy of the Safety Data Sheet (SDS), contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

HOW SUPPLIED

KetoMed (ketoprofen) Solution 100 mg/mL is available in 50 mL and 100 mL multidose bottles.

Store at 20°C - 25°C (68°F - 77°F); excursions permitted between 15°C - 30°C (59°F - 86°F). Do not freeze.

Use within 28 days of first puncture.

DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS DERIVED FROM THE USE OF SUCH PRODUCTS

Any unused product or waste materials should be disposed of in accordance with local requirements.

Approved by FDA under ANADA # 200-625

KetoMed™ is a trademark of Bimeda, Inc.

MANUFACTURED FOR:

Bimeda, Inc.

Le Sueur, MN 56058

www.bimeda.com

N.A. CORP. ADDRESS:

Bimeda, Inc.

One Tower Lane, Oakbrook Terrace, IL 60181

For further information and queries, please contact Bimeda at (888) 524-6332 or US-Info@Bimeda.com.

www.BimedaUS.com

